



## Role of oxidants in male infertility: rationale, significance, and treatment

Ashok Agarwal, PhD\*, Ramadan A. Saleh, MD

*Center for Advanced Research in Human Reproduction, Infertility, and Sexual Function;  
Urological Institute, and Department of Obstetrics/Gynecology, The Cleveland Clinic Foundation,  
Desk A 19.1, 9500 Euclid Avenue, Cleveland, OH 44195, USA*

Recently, the generation of oxidants, also described as reactive oxygen species (ROS), in the male reproductive tract has become a real concern because of their potential toxic effects, at high levels, on sperm quality and function [1,2]. ROS are highly reactive oxidizing agents belonging to the class of free radicals. As is true for all cells living under aerobic conditions, spermatozoa constantly face the oxygen (O<sub>2</sub>) paradox. O<sub>2</sub> is required to support life, but its metabolites, such as ROS, can modify cell functions, endanger cell survival, or both [3]; hence, ROS must be inactivated continuously to maintain only the small amount necessary to maintain normal cell function. A battery of different antioxidants normally protects against oxidants [4]. Oxidative stress develops when oxidants outnumber antioxidants, peroxidation products develop, and these phenomena cause pathologic effects [5,6].

In the context of human reproduction, a balance normally exists between ROS production and antioxidant scavenging activities in the male reproductive tract. As a result of such balance, only minimal amounts of ROS remain, and they are needed for the regulation of normal sperm functions, such as sperm capacitation, the acrosome reaction, and sperm–oocyte fusion [7,8]. The production of excessive amounts of ROS in semen can overwhelm the antioxidant defense mechanisms of spermatozoa and seminal plasma and causes oxidative stress [1,2,5].

Recent reports indicate that high levels of ROS are detected in the semen of 25% to 40% of infertile men [3,9]. Spermatozoa are particularly susceptible to the damage induced by excessive ROS because their plasma membranes contain large quantities of polyunsaturated fatty acids (PUFA) [10], and their cytoplasm contains low concentrations of scavenging enzymes [3]. In addition, the intracellular antioxidant enzymes cannot protect the plasma membrane that surrounds the acrosome and the tail, forcing spermatozoa to supplement their limited intrinsic antioxidant defenses by depending on the protection afforded by the seminal plasma [11,12].

Production of high levels of ROS in the reproductive tract is detrimental not only to the fluidity of the sperm plasma membrane but also to the integrity of DNA in the sperm nucleus [7]. Strong evidence suggests that DNA fragmentation commonly observed in the spermatozoa of infertile men is mediated by high levels of ROS [13,14].

This review discusses the mechanisms by which oxidants develop in semen and their role in physiology and pathology of sperm. Topics include the clinical implications of the presence of high levels of seminal ROS to the fertility potential of infertile men. In addition, research is summarized on treatment strategies for infertile men in whom oxidative stress has a significant role.

### **Mechanisms of oxidant generation in human semen**

A variety of semen components, including morphologically abnormal spermatozoa, precursor

---

\* Corresponding author.

*E-mail address:* agarwaa@ccf.org (A. Agarwal).

<http://www.clevelandclinic.org/reproductiveresearchcenter>

germ cells, and leukocytes, are capable of generating ROS. Seminal leukocytes and morphologically abnormal spermatozoa are the main sources of ROS in human ejaculates [15].

*Reactive oxygen species production by spermatozoa*

Clear evidence suggests that human spermatozoa can produce ROS [16–18]. Levels of ROS production by sperm correlate negatively with quality of sperm in the original semen [19]. The link between poor sperm quality and increased ROS generation lies in the retention of excess residual cytoplasm (cytoplasmic droplets) in abnormal spermatozoa. When spermatogenesis is impaired, the cytoplasmic extrusion mechanisms are defective. Spermatozoa released from the germinal epithelium carrying surplus residual cytoplasm are thought to be immature and functionally defective [20]. Retention of residual cytoplasm by spermatozoa is positively correlated with ROS generation via mechanisms that may be mediated by the cytosolic enzyme glucose-6-phosphate-dehydrogenase (G<sub>6</sub>PD). This enzyme controls the rate of glucose flux through the hexose monophosphate shunt, which, in turn, controls the intracellular availability of nicotinamide adenine dinucleotide phosphate (NADPH) (Fig. 1). The latter is used as a source of electrons by spermatozoa to fuel the generation of ROS by an enzyme system known as NADPH-oxidase [21].

Spermatozoa may generate ROS in two ways: (1) as a result of the NADPH-oxidase system at the level of the sperm plasma membrane, and (2) as a result of the NADH-dependent oxidoreductase (diphorase) at the level of mitochondria [16]. The mitochondrial system is the major source of ROS in spermatozoa from infertile men [22]. The primary ROS generated in human spermatozoa is the superoxide anion (O<sub>2</sub><sup>-</sup>). This one-electron reduction product of O<sub>2</sub> secondarily reacts with itself in a dismutation reaction, which is greatly accelerated by superoxide dismutase, to generate hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). In the presence of transition metals such as iron and copper, H<sub>2</sub>O<sub>2</sub> and O<sub>2</sub><sup>-</sup> can interact to generate the extremely pernicious hydroxyl radical (OH) (Haber-Weiss reaction) as shown in the following equation: O<sub>2</sub><sup>-</sup> + H<sub>2</sub>O<sub>2</sub> → OH + OH<sup>-</sup> + O<sub>2</sub>.

Alternatively, the hydroxyl radical can be produced from hydrogen peroxide (Fenton reaction), which requires a reducing agent such as ascorbate or ferrous ions, as shown in the equation: H<sub>2</sub>O<sub>2</sub> + Fe<sup>-2+</sup> → Fe<sup>3+</sup> + OH + OH<sup>-</sup>. The hydroxyl radical is thought to be an extremely powerful initiator of the lipid peroxidation cascade and can precipitate loss of sperm functions.

A study performed at the authors' center indicated that sperm production of ROS is significantly increased by the repeated cycles of centrifugation involved in the sperm preparation techniques used for assisted conception [23]. The duration of centrifugation was found to be more




Fig. 1. Mechanism of ROS production by immature spermatozoa (spermatozoa with excess residual cytoplasm). H<sub>2</sub>O<sub>2</sub>, hydrogen peroxide; NADPH, nicotinamide adenine dinucleotide phosphate; O<sub>2</sub><sup>-</sup>, superoxide anion.

important than the force of centrifugation in inducing sperm production of ROS [24]. Sperm production of ROS increases with the increase of sperm concentration and decreases with time [25]. Results from recent studies indicate a significant variation in ROS production in subsets of human spermatozoa at different stages of maturation [17,26]. Following isolate gradient fractionation of ejaculated sperm, ROS production was found to be highest in immature sperm with abnormal head morphology and cytoplasmic retention and lowest in mature sperm and immature germ cells. The levels of ROS production by immature sperm were directly correlated with sperm nuclear DNA damage values in mature sperm and inversely correlated with the recovery of motile mature sperm. These findings led to the hypothesis that oxidative damage of mature sperm by ROS-producing immature sperm during their comigration from seminiferous tubules to the epididymis may be an important cause of male infertility.

#### *Reactive oxygen species production by leukocytes*

Peroxidase-positive leukocytes are the major source of ROS in semen [27–29]. Peroxidase-positive leukocytes include polymorphonuclear leukocytes, which represent 50% to 60% of all seminal leukocytes, and macrophages, which represent 20% to 30% of all seminal leukocytes [30,31]. Peroxidase-positive leukocytes in semen are contributed largely by the prostate and seminal vesicles [32]. The capacity of leukocytes to generate ROS depends on their activation, which may occur in response to a variety of stimuli, including inflammation and infection [33]. During activation, NADPH production is increased, and the myeloperoxidase system of leukocytes is activated, leading to a respiratory burst with subsequent release of high levels of ROS [34]. Such an oxidative burst is thought to be an effective early defense that kills the microbes in cases of infection [35].

Sperm damage from ROS produced by leukocytes may occur when seminal plasma is removed during sperm preparation for assisted reproduction [28], or when seminal leukocyte concentrations are abnormally high, such as in leukocytospermia [29]. Sperm damage from leukocyte-derived ROS may happen even at leukocyte concentrations below the World Health Organization's cutoff value for leukocytospermia, that is, greater than  $1 \times 10^6$  peroxidase-positive leukocytes/mL of semen [36]. In addition, results

from the most recent studies performed at the authors' center indicate that seminal leukocytes may have a role in stimulating ROS production by human spermatozoa [37]. The exact mechanisms underlying such stimulation are not clear and may involve direct sperm–leukocyte contact or may be mediated by soluble products released by the leukocytes.

#### **Reactive oxygen species and sperm physiology**

Limited amounts of ROS can intervene physiologically in the regulation of some sperm functions [38]. It has been observed that the addition of low amounts of ROS to human spermatozoa enhances their ability to bind zona pellucida, an effect that is reversed by the addition of vitamin E. More recently, the incubation of spermatozoa with low concentrations of  $H_2O_2$  was found to stimulate sperm capacitation, hyperactivation, and the ability of spermatozoa to undergo the acrosome reaction and oocyte fusion [3,39]. ROS other than  $H_2O_2$ , such as nitric oxide and  $O_2^-$ , were also found to promote sperm capacitation and the acrosome reaction [40]. Sperm capacitation is a priming process that prepares spermatozoa in vivo in such a way that they reach the site of fertilization at the appropriate time and rapidly undergo the acrosome reaction upon contacting the zona pellucida. During the initial stages of capacitation, the intracellular concentration of calcium starts to rise, ROS generation is initiated, cyclic adenosine monophosphate (cAMP) concentrations increase, and spermatozoa develop a highly vigorous form of motility known as hyperactivation. When spermatozoa reach the oocyte, they fall under the influence of progesterone, which induces a sudden influx of extracellular calcium into the acrosomal region of the sperm head. As a result, spermatozoa become highly sensitized and ready to undergo the acrosome reaction [39].

#### **Reactive oxygen species and sperm toxicity**

Virtually every human ejaculate is contaminated with potential sources of ROS, such as peroxidase-positive leukocytes and morphologically abnormal spermatozoa. It follows that some of the sperm cells in every ejaculate incur oxidative damage and a concomitant loss of function. The extent of damage caused by ROS depends not only on the nature and amount of ROS involved but also on the moment and duration of ROS

exposure and the extracellular factors, such as temperature, oxygen tension, and the composition of the surrounding environment, including ions, proteins, and ROS scavengers.

#### *Lipid peroxidation of sperm plasma membrane*

Lipid peroxidation can be defined broadly as “oxidative deterioration of PUFA: ie, fatty acids that contain more than two carbon–carbon double bonds” [41]. Lipid peroxidation attacks the fluidity of sperm plasma membrane, with subsequent loss of the ability for oocyte fusion. The propagation of lipid peroxidation through sperm populations depends on the antioxidant strategies employed by spermatozoa and seminal plasma. One of the by-products of lipid peroxidation is malondialdehyde, which has been used as an end product in biochemical assays to monitor the degree of peroxidative damage to spermatozoa [38]. This assay correlates closely with the degree to which sperm motility and the capacity for oocyte fusion are impaired [42,43].

#### *Impairment of sperm motility*

Increased formation of ROS has been correlated with reduction of sperm motility [11,44]. The link between ROS and reduced motility may be explained by a cascade of events that result in a decrease in axonemal protein phosphorylation and sperm immobilization [45]. Another hypothesis is that  $H_2O_2$  can diffuse across the membranes into the cells and inhibit the activity of enzymes such as  $G_6PDH$ , leading to a decrease in the availability of NADPH and a concomitant accumulation of oxidized glutathione and reduced glutathione. These changes can cause a decrease in the antioxidant defenses of the spermatozoa, which ultimately leads to the peroxidation of membrane phospholipids [46].

#### *Sperm DNA damage*

Two factors protect spermatozoal DNA from oxidative insult: the characteristic tight packaging of sperm DNA and the antioxidants in seminal plasma [47]. Exposing the sperm to artificially produced ROS causes DNA damage in the form of modification of all bases, production of base-free sites, deletions, frame shifts, DNA cross-links, and chromosomal rearrangements [48]. Oxidative stress is also associated with high frequencies of single and double DNA strand breaks [14]. This information may have important clinical implications, particularly in the context of assisted reproductive

techniques (ART). Spermatozoa selected for ART usually originate from an environment experiencing oxidative stress, and a high percentage of these sperm may have damaged DNA [13,49]. There is a substantial risk that spermatozoa carrying damaged DNA are used in this form of therapy [50]. When intrauterine insemination (IUI) or in vitro fertilization (IVF) is used, such damage may not be a cause of concern because the collateral peroxidative damage to the sperm plasma membrane ensures that fertilization cannot occur with a DNA-damaged sperm. When intracytoplasmic sperm injection (ICSI) is used, this natural selection barrier is bypassed, and a spermatozoon with damaged DNA is directly injected into the oocyte [7,50].

Whether DNA-damaged spermatozoa used in ICSI can impair the process of fertilization and embryo development is not clear. A recent study indicated that spermatozoa with significantly damaged DNA retain the capacity for fertilization [7]; however, the percentage of sperm with DNA damage negatively correlates with fertilization rates [14,49]. In addition, a recent study linked sperm DNA damage to increased rates of early embryo death [51].

#### **Antioxidant defense mechanisms**

The seminal plasma contains enzymatic antioxidants, such as superoxide dismutase, the glutathione peroxidase/glutathione reductase system, and catalase [52], as well as nonenzymatic antioxidants, such as ascorbate (vitamin C) [53],  $\alpha$ -tocopherol (vitamin E) [54], taurine, and hypotaurine [55]. The seminal plasma of fertile men has a higher total antioxidant capacity (TAC) than the seminal plasma of infertile men [56]. Nevertheless, the pathologic levels of ROS detected in the semen of infertile men are more likely caused by increased ROS production than by reduced antioxidant capacity of the seminal plasma [12].

Antioxidant defense mechanisms include three levels of protection: prevention, interception, and repair. Prevention of ROS formation is the first line of defense against an oxidative insult. An example is the binding of metal ions, iron and copper ions in particular, which prevents them from initiating a chain reaction [4]. When transition metals become loosely bound to ROS, they can produce more reactive oxidants, particularly  $OH^\bullet$  [57]. Free radicals have a tendency toward chain reaction, that is, a compound carrying an unpaired electron will react with another compound to generate an unpaired electron as “radical begets radi-

cal.” The basic problem is to intercept a damaging species from further activity, which is the process of deactivation leading to the formation of non-radical end products [4].  $\alpha$ -Tocopherol, a chain-breaking antioxidant, inhibits lipid peroxidation in membranes by scavenging peroxy (RO $\cdot$ ) and alkoxy (ROO $\cdot$ ) radicals, thereby breaking the chain reaction. The ability of  $\alpha$ -tocopherol to maintain a steady-state rate of peroxy radical reduction in the plasma membrane depends on the recycling of  $\alpha$ -tocopherol by external reducing agents such as ascorbate or thiols. In this way,  $\alpha$ -tocopherol is able to function again as a free radical chain-breaking antioxidant, even though its concentration is low [58]. Protection from the effects of oxidants also can be brought about by repairing the damage once it has occurred. Unfortunately, spermatozoa are unable to repair the damage induced by oxidative stress, because they lack the cytoplasmic enzyme systems required to accomplish the repair [38,52].

## Methods to assess seminal oxidative stress

### Reactive oxygen species–total antioxidant capacity score

Because oxidative stress is caused by an imbalance between levels of ROS production and antioxidant protection in semen, it is conceivable that assessment of seminal oxidative stress will rely on measurement of ROS as well as TAC of semen. Levels of seminal ROS can be measured accurately by a chemiluminescence assay [59], and TAC of the seminal plasma can be assessed by an enhanced chemiluminescence assay [60]. The fact that neither ROS nor TAC alone can adequately quantify seminal oxidative stress led to the logical conclusion that combining these two variables may be a

better index for diagnosis of the overall oxidative stress affecting spermatozoa. This conclusion was behind the development of a statistical formula described as the ROS-TAC score for assessment of seminal oxidative stress in infertile men [61]. This score minimizes the variability of the individual parameters of oxidative stress (ROS alone or TAC alone). The ROS-TAC score was based on a group of normal healthy fertile men who had very low levels of ROS. The composite ROS-TAC score is representative of the fertile population, and scores significantly lower (<30) indicate infertility [61].

### Measurement of reactive oxygen species in neat semen

More recently, the authors' group introduced an additional test for accurate and reliable assessment of oxidative stress by measurement of ROS levels directly in neat (raw) semen using a chemiluminescence assay [62]. The test was subjected to quality control studies and proved to be an accurate measure for seminal oxidative stress status. The maximum ROS level observed in neat semen from normal healthy donors, with normal genital examination and normal standard semen parameters, was  $1.5 \times 10^4$  cpm/20 million sperm/mL. Infertile men were reliably classified into oxidative stress–positive ( $\geq 1.5 \times 10^4$  cpm/20 million sperm/mL) or oxidative stress–negative ( $< 1.5 \times 10^4$  cpm/20 million sperm/mL), regardless of their clinical diagnosis or the results of standard semen analysis. Assessment of ROS directly in neat semen had diagnostic and prognostic capabilities identical to those based on the ROS-TAC score (Table 1) [63]. Levels of ROS in neat semen were strongly correlated with levels of ROS in washed semen and with ROS-TAC scores.

Table 1

Median (25th and 75th percentile) values of reactive oxygen species (ROS) in neat semen, ROS in washed semen, total antioxidant capacity (TAC) in seminal plasma, and ROS-TAC score in donors, oxidative stress (OS)–negative, and OS-positive patients

Variable	Donors (n = 9)	OS-negative (n = 11)	OS-positive (n = 23)	A	B	C
ROS in neat semen ( $\times 10^4$ cpm)	0.3 (0.2, 0.9)	0.3 (0.2, 1)	19 (6, 143)	0.9	0.0001	0.0001
ROS in washed semen ( $\times 10^4$ cpm)	10 (4, 17)	79 (13, 177)	1468 (514, 11496)	0.03	0.0003	0.0008
TAC (Trolox equivalent)	908 (736, 1129)	797 (575, 966)	739 (627, 1047)	0.31	0.24	0.94
ROS-TAC score	53 (51, 55)	49 (47, 52)	36 (32, 44)	0.4	0.001	0.001

Results were analyzed by Wilcoxon rank-sum test;  $P < 0.05$  was significant.

Abbreviations: A,  $P$  value of donors versus OS-negative patients; B,  $P$  value between donors versus OS-positive patients; C, OS-negative patients versus OS-positive patients.

### Clinical significance of seminal oxidative stress testing

Accurate assessment of seminal levels of oxidative stress may help to diagnose infertility cases in which oxidative stress has a major role (Table 2). Levels of ROS were found to be increased significantly in men with spinal cord injury and were associated with poor sperm motility and morphology [9]. A similar increase of seminal ROS was observed in infertile men with varicoceles [18]. Patients with varicocele were also found to have low levels of TAC in their seminal plasma and might benefit from antioxidant supplementation [18]. In a recent study, varicocelectomy resulted in a significant increase in pregnancy and live birth rates for couples who underwent IUI, although standard semen parameters were not improved in all patients [64]. Such improvement in pregnancy rates following varicocelectomy may result from the correction of a functional factor not tested during routine semen analysis, such as seminal oxidative stress. The new measures of seminal oxidative stress, such as the ROS-TAC score and ROS levels in neat semen, are tools for successful discrimination between fertile and infertile men [37,61]. Infertile men with chronic prostatitis or prostatodynia were found to have lower ROS-TAC scores than normal sperm donors [65]. Similarly, infertile men with male-factor or idiopathic diagnoses had significantly lower ROS-TAC scores than normal sperm donors. In addition, infertile men with male-factor diagnoses who eventually were able to initiate a successful pregnancy had significantly higher ROS-TAC scores than men who failed to initiate a pregnancy [66]. The ROS-TAC score

may serve as an important measure in identifying men with a clinical diagnosis of infertility who are likely to initiate a pregnancy over time [33]. The average ROS-TAC score for fertile men who had undergone vasectomy was nearly identical to that of fertile donors [60].

### Treatment strategies

Given the major role of oxidative stress in the pathogenesis of male infertility, treatment strategies with the goal of reducing levels of seminal oxidative stress are necessary for natural and assisted conception.

#### *Elimination of reactive oxygen species sources from semen*

Once seminal oxidative stress is diagnosed, treatment plans should focus on identification and elimination of the sources of enhanced ROS production in semen. The differentiation between spermatozoal and leukocyte sources of ROS may significantly affect therapeutic strategies for infertility cases in which oxidative stress has a causative role. Seminal leukocytes should be considered potentially detrimental and must be monitored carefully. The underlying etiologic factors for abnormal leukocyte infiltration into semen (leukocytospermia), such as inflammation, infection, and cigarette smoking, should be determined and properly addressed. The interaction between ROS-producing cells in semen (leukocytes or abnormal spermatozoa) and normal spermatozoa must be minimized. Sperm preparation techniques used for ART, such as density gradient and swim-

Table 2

Mean and standard deviation between reactive oxygen species (ROS), total antioxidant capacity (TAC), and ROS-TAC score in subgroups of infertility patients and controls

Diagnosis	ROS Log (ROS+1)	<i>P</i> value versus controls*	TAC (Trolox equivalent)	<i>P</i> value versus controls*	ROS-TAC score	<i>P</i> value versus controls*
Control (n = 24)	1.39 ± 0.73	—	1650.93 ± 532.22	—	50.00 ± 10.00	—
Varicocele (n = 55)	2.10 ± 1.21	0.02	1100.11 ± 410.30	0.0002	34.87 ± 13.54	0.0001
Varicocele with prostatitis (n = 8)	3.25 ± 0.89	0.0002	1061.42 ± 425.11	0.03	22.39 ± 13.48	0.0001
Vasectomy reversal (infertile, n = 23)	2.65 ± 1.01	0.0004	1389.89 ± 723.92	0.30	33.22 ± 15.24	0.0002
Vasectomy reversal (fertile, n = 12)	1.76 ± 0.86	0.80	1876.93 ± 750.82	0.62	49.35 ± 12.25	1.00
Idiopathic infertility (n = 28)	2.29 ± 1.20	0.01	1051.98 ± 380.88	0.0003	32.25 ± 14.40	0.0001

\* Pairwise *P* values from Student's *t*-test adjusted using Dunnett's method.

up methods, can be used for separation of spermatozoa with good quality that may have a potential to fertilize; however, clinicians should be aware that the repeated cycles of centrifugation routinely performed in these procedures may induce ROS generation by spermatozoa. Such increases in ROS levels together with lack of antioxidant protection owing to removal of seminal plasma may expose the spermatozoa to risk of oxidative damage. The use of sperm preparation techniques with shorter centrifugation periods may reduce the risk of oxidative stress-induced injury to sperm.

#### *Antioxidant supplementation*

It has been postulated that protective agents against ROS, such as antioxidants, may be useful therapeutic agents for male infertility [67]; however, the results of clinical trials in which antioxidants have been used for the treatment of infertile men are controversial [68]. The results of 6 months of supplementation with vitamin E indicated an improvement in sperm motility limited to the patients whose original sperm motility was greater than 15% [69]. A combination of vitamin E and C administered orally for 2 months resulted in improvement of sperm concentration [13]. Administration of glutathione via intramuscular injection for 2 months resulted in improvement of sperm concentration, motility, and morphology [70]. To the contrary, some studies have found no changes in sperm characteristics following treatment with vitamin C or E [71–73]. In a recent prospective clinical trial, the treatment of infertile men with a combination of vitamins A plus E and essential fatty acids improved the sperm concentration in oligozoospermic men but did not improve sperm motility or morphology [74].

Levels of seminal ROS and TAC were not predetermined in these studies before antioxidant supplementation. Accurate assessment of levels of seminal ROS and TAC may help to identify subgroups of infertility patients with high ROS, low TAC, or both who may benefit from antioxidant supplementation.

The results of *in vitro* trials using antioxidants are certainly not better than the results of *in vivo* trials, and the potential advantages of antioxidants in assisted reproduction are the subject of major debate [75,76]. During sperm preparation for ART, spermatozoa are washed free from their seminal plasma and left vulnerable to oxidative stress owing to a lack of antioxidant protection. It has been speculated that supplementing the

sperm preparation media with antioxidants may provide some degree of protection to the spermatozoa against oxidative stress-induced damage [77]. On this basis, several studies have examined the potential role of antioxidant supplementation *in vitro* [78]. Vitamin E has been shown to inhibit sperm lipid peroxidation *in vitro* [79], to enhance the ability of spermatozoa to fuse with zona-free hamster oocytes [38], and to improve zona pellucida binding [80]. Supplementing the sperm preparation media with a combination of vitamins C and E was associated with decreased ROS production by the sperm [81]. In another study, supplementation with superoxide dismutase was associated with improved rates of acrosome reaction and preservation of sperm motility [8]. Baker et al (1996) have shown that glutathione, hypotaurine, and catalase can reduce the loss of motility caused by leukocyte-generated ROS. Lewis et al [82] recommended adding low-molecular weight antioxidants such as ascorbate to sperm during preparation for ART, especially in asthenozoospermic patients, on the basis of low levels of ascorbate in those patients. They also indicated that it would be of more clinical benefit to add these antioxidants directly to the sperm rather than using dietary supplements, because defective seminal vesicle function may be one of the causes of reduced ascorbate levels in these patients.

When vitamin E or C was added to the sperm preparation media during density gradient separation using Percoll, spermatozoa were protected from DNA damage [77]; however, when both of these vitamins were used together, DNA damage increased. Twigg et al [47] found that albumin can be an important means of neutralizing lipid peroxide-mediated damage to the sperm plasma membrane and DNA.

Sperm motion characteristics were reduced significantly when a combination of vitamins E and C was used, with the greatest reduction observed with the highest concentrations [81]. It was concluded that supplementing the sperm preparation media with vitamins E and C was not beneficial.

No definitive conclusion has been reached, with some investigators overestimating and others underestimating the potential benefits of antioxidants in the treatment of male infertility.

#### **Future directions**

Future research efforts should be directed toward understanding why spermatozoa of some patients become overreactive in the generation of

ROS. Insight into the molecular basis of these defects is vital to identify the cause of unexplained male infertility. If the error in spermatogenesis that leads to such atypical activity (excessive ROS production) could be defined, it would provide a rational basis for the design of effective therapies. The period of sperm differentiation during which this self-destructive activity first appears must be determined. Further efforts are required to identify the sperm population at risk of collateral peroxidative damage to the sperm membrane. An interesting area of future research is to investigate precisely the oxidative damage to sperm DNA and its effects on male fertility potential and the outcome of ART.

### Summary

Interest in the physiologic and pathologic effects of ROS on male fertility is growing. The controlled generation of very low amounts of ROS seems to regulate the acquisition of sperm-fertilizing ability. High levels of ROS endanger sperm function and viability. Oxidative stress can arise as a consequence of excessive production of ROS or impaired antioxidant defense mechanisms in semen. Oxidative stress precipitates a range of pathologies that are thought to affect the male reproductive system. ROS-mediated peroxidative damage to the sperm plasma membrane may account for the defective sperm functions observed in a high proportion of infertility patients. High levels of ROS in semen have been correlated with reduced sperm motility and damage to sperm nuclear DNA.

Determining the levels and the sources of excessive ROS production in human semen and precise evaluation of the scavenger system may be useful tools to develop therapeutic strategies for male infertility. Therapeutic interventions directed toward the isolation of mature spermatozoa by *in vitro* separation techniques may benefit infertile men in whom oxidative stress has a significant role. Nevertheless, this approach may be limited by the fact that the prolonged centrifugation involved in these procedures may enhance ROS production by spermatozoa. Clinical trials using antioxidants *in vivo* and *in vitro* have resulted in a major debate, and further research is required before one can be optimistic about a role for antioxidants in the treatment of infertile men. A double-blind, placebo-controlled study underway at the authors' center is evaluating the effects of treating infertile

men with a combination of vitamins E and C after adjusting for all causes of controversy.

### References

- [1] Sharma RK, Agarwal A. Role of reactive oxygen species in male infertility. *Urology* 1996;48:835–50.
- [2] Sikka SC, Rajasekaran M, Hellstrom WJG. Role of oxidative stress and antioxidants in male infertility. *J Androl* 1995;16:464–8.
- [3] de Lamirande E, Gagnon C. Impact of reactive oxygen species on spermatozoa: a balancing act between beneficial and detrimental effects. *Hum Reprod* 1995;10:15–21.
- [4] Sies H. Strategies of antioxidant defense. *Eur J Biochem* 1993;215:213–9.
- [5] Sikka SC. Relative impact of oxidative stress on male reproductive function. *Curr Med Chem* 2001; 8:851–62.
- [6] Spitteler G. Review: on the chemistry of oxidative stress. *J Lipid Mediat* 1993;7:77–82.
- [7] Aitken RJ. The Amoroso lecture: the human spermatozoon—a cell in crisis? *J Reprod Fertil* 1999;115:1–7.
- [8] Griveau JF, Le Lannou D. Reactive oxygen species and human spermatozoa. *Int J Androl* 1997;20:61–9.
- [9] Padron OF, Brackett NL, Sharma RK, et al. Seminal reactive oxygen species, sperm motility and morphology in men with spinal cord injury. *Fertil Steril* 1997;67:1115–20.
- [10] Alvarez JG, Storey BT. Differential incorporation of fatty acids into and peroxidative loss of fatty acids from phospholipids of human spermatozoa. *Mol Reprod Dev* 1995;42:334–46.
- [11] Iwasaki A, Gagnon C. Formation of reactive oxygen species in spermatozoa of infertile patients. *Fertil Steril* 1992;57:409–16.
- [12] Zini A, de Lamirande E, Gagnon C. Reactive oxygen species in the semen of infertile patients: levels of superoxide dismutase- and catalase-like activities in seminal plasma. *Int J Androl* 1993;16:183–8.
- [13] Kodama H, Yamaguchi R, Fukuda J, et al. Increased oxidative deoxyribonucleic acid damage in the spermatozoa of infertile male patients. *Fertil Steril* 1997;65:519–24.
- [14] Sun JG, Jurisicova A, Casper RF. Deletion of deoxyribonucleic acid fragmentation in human sperm: correlation with fertilization *in vitro*. *Biol Reprod* 1997;56:602–7.
- [15] Kessopoulou E, Tomlinson MJ, Banat CLR, et al. Origin of reactive oxygen species in human semen-spermatozoa or leukocytes. *J Reprod Fertil* 1992; 94:463–70.
- [16] Aitken RJ, Buckingham DW, West KM. Reactive oxygen species and human spermatozoa: analysis of the cellular mechanisms involved in luminol- and lucigenin-dependent chemiluminescence. *J Cell Physiol* 1992;151:466–77.

- [17] Gil-Guzman E, Ollero M, Lopez MC, et al. Differential production of reactive oxygen species by subsets of human spermatozoa at different stages of maturation. *Hum Reprod* 2001;16:1922–30.
- [18] Hendin B, Kolettis P, Sharma RK, et al. Varicocele is associated with elevated spermatozoal reactive oxygen species production and diminished seminal plasma antioxidant capacity. *J Urol* 1999;161:1831–4.
- [19] Gomez E, Irvine DS, Aitken RJ. Evaluation of a spectrophotometric assay for the measurement of malondialdehyde and 4-hydroxyalkenals in human spermatozoa: relationships with semen quality and sperm function. *Int J Androl* 1998;21:81–94.
- [20] Huszar G, Sbracia M, Vigue L, et al. Sperm plasma membrane remodeling during spermiogenic maturation in men: relationship among plasma membrane beta 1,4-galactosyltransferase, cytoplasmic creatine phosphokinase and creatine phosphokinase isoform ratios. *Biol Reprod* 1997;56:1020–4.
- [21] Aitken RJ, Fisher HM, Fulton N, et al. Reactive oxygen species generation by human spermatozoa is induced by exogenous NADPH and inhibited by flavoprotein inhibitors diphenylene iodonium and quinacrine. *Mol Reprod Dev* 1997;47:468–82.
- [22] Plante M, de Lamirande E, Gagnon C. Reactive oxygen species released by activated neutrophils, but not by deficient spermatozoa, are sufficient to affect normal sperm motility. *Fertil Steril* 1994;62:387–93.
- [23] Agarwal A, Ikemoto I, Loughlin KR. Effect of sperm washing on reactive oxygen species level in semen. *Arch Androl* 1994;33:157–62.
- [24] Shekarriz M, Thomas Jr AJ, Agarwal A. A method of human semen centrifugation to minimize the iatrogenic sperm injuries caused by reactive oxygen species. *Eur Urol* 1995;28:31–5.
- [25] Shekarriz M, Thomas Jr AJ, Agarwal A. Incidence and level of seminal reactive oxygen species in normal men. *Urology* 1995;45:103–7.
- [26] Ollero M, Gil-Guzman E, Lopez MC, et al. Characterization of subsets of human spermatozoa at different stages of maturation: implications in the diagnosis and treatment of male infertility. *Hum Reprod* 2001;16:1912–21.
- [27] Aitken RJ, West KM. Analysis of the relationship between reactive oxygen species production and leukocyte infiltration in fractions of human semen separated on Percoll gradients. *Int J Androl* 1990;3:433–51.
- [28] Ochsendorf FR. Infections in the male genital tract and reactive oxygen species. *Hum Reprod* 1999;5:399–420.
- [29] Shekarriz M, Sharma RK, Thomas Jr AJ, et al. Positive myeloperoxidase staining (Endtz Test) as an indicator of excessive reactive oxygen species formation in semen. *J Assist Reprod Genet* 1995;12:70–4.
- [30] Fedder J, Askjaer SA, Hjort T. Non-spermatozoal cells in semen: relationship to other semen parameters and fertility status of the couple. *Arch Androl* 1993;31:95–103.
- [31] Wolff H, Anderson DJ. Immunohistologic characterization and quantitation of leukocyte subpopulation in human semen. *Fertil Steril* 1988;49:497–503.
- [32] Wolff H. The biologic significance of white blood cells in semen. *Fertil Steril* 1995;63:1143–57.
- [33] Pasqualotto FF, Sharma RK, Nelson DR, et al. Relationship between oxidative stress, semen characteristics, and clinical diagnosis in men undergoing fertility investigation. *Fertil Steril* 2000;73:459–64.
- [34] Blake DR, Allen RE, Lunec J. Free radical in biological systems—a review oriented to inflammatory processes. *Br Med Bull* 1987;43:371–85.
- [35] Saran M, Beck-Speier I, Fellerhoff B, et al. Phagocytic killing of microorganisms by radical processes: consequences of the reaction of hydroxyl radicals with chloride yielding chlorine atoms. *Free Radic Biol Med* 1999;26:482–90.
- [36] Sharma RK, Pasqualotto FF, Nelson DR, et al. Relationship between seminal white blood cell counts and oxidative stress in men treated at an infertility clinic. *J Androl* 2001;22:575–83.
- [37] Saleh RA, Agarwal A, Kandirali E, et al. Leukocytospermia is associated with increased reactive oxygen species production by human spermatozoa. *Fertil Steril* 2002 (in press).
- [38] Aitken RJ, Clarkson JS, Fishel S. Generation of reactive oxygen species, lipid peroxidation, and human sperm function. *Biol Reprod* 1989;40:183–97.
- [39] Aitken RJ, Paterson M, Fisher H, et al. Redox regulation of tyrosine phosphorylation in human spermatozoa is involved in the control of human sperm function. *J Cell Sci* 1995;108:2017–25.
- [40] Zini A, de Lamirande E, Gagnon C. Low levels of nitric oxide promote human sperm capacitation in vitro. *J Androl* 1996;16:424–31.
- [41] Halliwell B. Tell me about free radicals, doctor: a review. *J R Soc Med* 1984;82:747–52.
- [42] Aitken RJ, Harkiss D, Buckingham D. Relationship between iron-catalyzed lipid peroxidation potential and human sperm function. *J Reprod Fertil* 1993;98:257–65.
- [43] Sidhu RS, Sharma RK, Thomas Jr AJ, et al. Relationship between creatine kinase activity and semen characteristics in subfertile men. *Int J Fertil Womens Med* 1998;43:192–7.
- [44] Agarwal A, Ikemoto I, Loughlin KR. Relationship of sperm parameters to levels of reactive oxygen species in semen specimens. *J Urol* 1994;152:107–10.
- [45] de Lamirande E, Gagnon C. Reactive oxygen species and human spermatozoa. II. Depletion of adenosine triphosphate (ATP) plays an important role in the inhibition of sperm motility. *J Androl* 1992;13:379–86.

- [46] Griveau JF, Dumont E, Renard B, et al. Reactive oxygen species, lipid peroxidation and enzymatic defense systems in human spermatozoa. *J Reprod Fertil* 1995;103:17–26.
- [47] Twigg J, Irvine DS, Houston P, et al. Iatrogenic DNA damage induced in human spermatozoa during sperm preparation: protective significance of seminal plasma. *Mol Hum Reprod* 1998;4:439–45.
- [48] Duru NK, Morshedi M, Oehninger S. Effects of hydrogen peroxide on DNA and plasma membrane integrity of human spermatozoa. *Fertil Steril* 2000;74:1200–7.
- [49] Lopes S, Jurisicova A, Sun J, et al. Reactive oxygen species: a potential cause for DNA fragmentation in human spermatozoa. *Hum Reprod* 1998;13:896–900.
- [50] Twigg J, Irvine DS, Aitken RJ. Oxidative damage to DNA in human spermatozoa does not preclude pronucleus formation at intracytoplasmic sperm injection. *Hum Reprod* 1998;13:1864–71.
- [51] Sakkas D, Mariethoz E, Manicardi G, et al. Origin of DNA damage in ejaculated human spermatozoa. *Rev Reprod* 1999;4:31–7.
- [52] Alvarez JG, Touchstone JC, Blasco L, et al. Spontaneous lipid peroxidation and production of hydrogen peroxide and superoxide in human spermatozoa: superoxide dismutase as major enzyme protectant against oxygen toxicity. *J Androl* 1987;8:336–48.
- [53] Fraga GG, Motchnik PA, Shigenaga MK. Ascorbic acid protects against endogenous oxidative DNA damage in human sperm. *Proc Natl Acad Sci USA* 1991;88:11003–6.
- [54] Moilanen J, Hovatta O, Lindroth L. Vitamin E levels in seminal plasma can be elevated by oral administration of vitamin E in infertile men. *Int J Androl* 1993;16:165–6.
- [55] Alvarez JG, Storey BT. Taurine, hypotaurine, epinephrine and albumin inhibit lipid peroxidation in rabbit spermatozoa and protect against loss of motility. *Biol Reprod* 1983;29:548–55.
- [56] Lewis SEM, Boyle PM, McKinney KA, et al. Total antioxidant capacity of seminal plasma is different in fertile and infertile men. *Fertil Steril* 1995;64:868–70.
- [57] Halliwell B. How to characterize a biological antioxidant. *Free Radic Res* 1990;9:1–32.
- [58] Buettner GR. The pecking order of free radicals and antioxidants, lipid peroxidation, alpha-tocopherol and ascorbate. *Arch Biochem Biophys* 1993;300:535–43.
- [59] Kobayashi H, Gil-Guzman E, Mahran AM, et al. Quality control of reactive oxygen species measurement by luminol-dependent chemiluminescence assay. *J Androl* 2000;22:568–74.
- [60] Kolettis P, Sharma RK, Pasqualotto FF, et al. The effects of seminal oxidative stress on fertility after vasectomy reversal. *Fertil Steril* 1999;71:249–55.
- [61] Sharma RK, Pasqualotto FF, Nelson DR, et al. The reactive oxygen species—total antioxidant capacity score is a new measure of oxidative stress to predict male infertility. *Hum Reprod* 1999;14:2801–7.
- [62] Saleh RA, Esfandiari N, Al-Dujaily S, et al. An accurate and reliable method for the diagnosis of seminal oxidative stress in infertile men [abstract O-274]. In: Programs and Abstracts of the 57th Annual Meeting of the American Society for Reproductive Medicine. Orlando (FL); 2001. p. 104.
- [63] Saleh RA, Esfandiari N, Sharma RK, et al. Diagnostic and prognostic value of measurement of reactive oxygen species in neat semen [abstract O-24]. In: Programs and Abstracts of the 57th Annual Meeting of the American Society for Reproductive Medicine. Orlando (FL); 2001. p. 9.
- [64] Daitch JA, Bedaiway MA, Pasqualotto FF. Varicocelectomy improves intrauterine insemination success rates among men with varicocele. *J Urol* 2001;165:1510–3.
- [65] Pasqualotto FF, Sharma RK, Agarwal A, et al. Seminal oxidative stress in chronic prostatitis patients. *Urology* 2000;55:881–5.
- [66] Pasqualotto FF, Sharma RK, Kobayashi H, et al. Oxidative stress in normospermic men undergoing infertility evaluation. *J Androl* 2001;73:459–64.
- [67] Rolf C, Cooper TG, Yeung CH, et al. Antioxidant treatment of patients with asthenozoospermia or moderate oligoasthenozoospermia with high-dose vitamin C and vitamin E: a randomized, placebo-controlled, double-blind study. *Hum Reprod* 1999;14:1028–33.
- [68] Martin-Du Pan RC, Sakkas D. Are antioxidants useful in the treatment of male infertility? *Hum Reprod* 1998;13:2984–5.
- [69] Suleiman SA, Ali ME, Zaki ZMS, et al. Lipid peroxidation and human sperm motility: protective role of vitamin E. *J Androl* 1996;17:530–7.
- [70] Lenzi A, Cualosso F, Gandini L, et al. Placebo-controlled, double-blind, cross-over trial of glutathione therapy in male infertility. *Hum Reprod* 1993;9:2044–50.
- [71] Abel BJ, Carswell G, Elton R, et al. Randomized trial of clomiphene citrate treatment and vitamin C for male infertility. *Br J Urol* 1982;54:780–4.
- [72] Geva E, Bartoov B, Zabludovsky N, et al. The effect of antioxidant treatment on human spermatozoa and fertilization rate in an in vitro fertilization program. *Fertil Steril* 1996;66:430–4.
- [73] Moilanen J, Hovatta O. Excretion of alpha-tocopherol into human seminal plasma after oral administration. *Andrologia* 1995;27:133–6.
- [74] Comhaire FH, Christophe AB, Zalata AA, et al. The effects of combined conventional treatment, oral antioxidants and essential fatty acids on sperm biology in subfertile men. *Prostaglandins Leukot Essent Fatty Acids* 2000;63:159–65.
- [75] Ford WCL, Whittington K. Antioxidant treatment for male subfertility: a promise that remains unfulfilled. *Hum Reprod* 1998;13:1416–9.

- [76] Tarin JJ, Brines J, Cano A. Is antioxidant therapy a promising strategy to improve human reproduction? *Hum Reprod* 1998;13:1415–24.
- [77] Hughes CM, Lewis SEM, McKelvey-Martin VJ, et al. The effects of antioxidant supplementation during Percoll preparation on human sperm DNA integrity. *Hum Reprod* 1998;13:1240–7.
- [78] Irvine DS. Glutathione as a treatment of male infertility. *Rev Reprod* 1996;1:6–12.
- [79] Aitken RJ, Clarkson JS. Significance of reactive oxygen species and antioxidants in defining the efficacy of sperm preparation techniques. *J Androl* 1988;9:367–76.
- [80] Kessopoulou E, Powers HJ, Sharma KK, et al. A double-blind randomized placebo cross-over controlled trial using the antioxidant vitamin E to treat reactive oxygen species associated male infertility. *Fertil Steril* 1995;64:825–31.
- [81] Donnelly ET, McClure N, Lewis S. Antioxidant supplementation in vitro does not improve human sperm motility. *Fertil Steril* 1999;72:484–95.
- [82] Lewis SEM, Sterling SEL, Young IS, et al. Comparison of individual antioxidants of sperm and seminal plasma in fertile and infertile men. *Fertil Steril* 1997;67:142–7.