

Effects of magnetic-activated cell sorting on sperm motility and cryosurvival rates

Tamer M. Said, M.D.,^a Sonja Grunewald, M.D.,^b Uwe Paasch, M.D., Ph.D.,^b Manja Rasch,^b Ashok Agarwal, Ph.D., H.C.L.D.,^a and Hans-Juergen Glander, M.D., Ph.D.^b

^aCenter for Advanced Research in Human Reproduction, Infertility and Sexual Function, Glickman Urological Institute and Department of Obstetrics-Gynecology, Cleveland Clinic Foundation, Cleveland, Ohio; and ^bDepartment of Dermatology/Andrology Unit, University of Leipzig, Leipzig, Germany

Objective: To evaluate the effect of magnetic-activated cell sorting in cryopreservation–thawing protocols on sperm motility and cryosurvival rate.

Design: Prospective-controlled study.

Setting: Andrology department at a university-based medical institution.

Patient(s): Ten healthy volunteer sperm donors.

Intervention(s): Sperm populations were separated using annexin-V magnetic-activated cell sorting before and after the cryopreservation–thawing process.

Main Outcome Measure(s): Sperm motility and cryosurvival rate.

Result(s): Annexin-negative sperm separated by magnetic-activated cell sorting had statistically significantly higher motility following cryopreservation–thawing than sperm that were not separated. Similarly, annexin-negative spermatozoa also had higher cryosurvival rate than sperm cryopreserved without magnetic-activated cell sorting and sperm that were annexin-positive.

Conclusion(s): Superparamagnetic annexin V-conjugated microbeads can separate spermatozoa with externalized phosphatidylserine, which is considered one of the early features of late apoptosis. The separation of a distinctive population of nonapoptotic spermatozoa with intact membranes may optimize the cryopreservation–thawing outcome. Magnetic-activated cell sorting using annexin-V microbeads enhances sperm motility and cryosurvival rates following cryopreservation. (*Fertil Steril*® 2005;83:1442–6. ©2005 by American Society for Reproductive Medicine.)

Key Words: Annexin, cryosurvival rate, magnetic-activated cell sorting, sperm

Cryopreservation of human semen is the most commonly accepted method of preserving male reproductive capacity. Cryopreserved spermatozoa may be used in assisted reproductive techniques (ART) (1), especially in cases where the patient elects to undergo vasectomy for contraception, or, most importantly, when a patient is diagnosed with cancer and the treatment may render him infertile (2). The indications for sperm cryobanking have been greatly expanded by recent breakthroughs in ART, in which immotile but viable sperm can be used successfully for oocyte fertilization through intracytoplasmic sperm injection (ICSI).

Despite recent methodological advances, cryopreservation exerts detrimental effects on spermatozoa that lead to significant decreases in sperm viability and motility and ultimately in decreased cryosurvival rates (CSR) (3). The fertility potential of cryopreserved mammalian spermatozoa is lower than that of fresh sperm. The reduction arises from both a lower post-thaw viability and sublethal dysfunction in

a proportion of the surviving subpopulation (4). Programmed cell death (apoptosis) most likely contributes to the decrease in sperm quality after cryopreservation (5).

The sperm plasma membrane is one of the key structures affected by cryopreservation that displays apoptotic features (6). Early phases of disturbed membrane functions are associated with asymmetry of the membrane phospholipids. The phospholipid phosphatidylserine (PS), which is normally present on the inner leaflet of the plasma membrane, becomes externalized to the outer leaflet (7). The externalization of PS is a known early marker for apoptosis (8). Because annexin-V has a high affinity for PS, it cannot pass through an intact sperm membrane. Therefore, when annexin-V binds to spermatozoa, it signifies that the integrity of the membrane has been disturbed (9).

Colloidal super-paramagnetic microbeads (~50 nm in diameter) conjugated with annexin-V may be used to separate dead and apoptotic spermatozoa by magnetic-activated cell sorting (MACS). Cells with PS that has externalized to the outer leaflet will bind to these microbeads. When placed into a column containing iron balls and passed through a strong magnetic field, those cells remain in the separation column. On the other hand, cells with intact membranes remain unlabeled and pass freely through the column (10, 11).

Received August 23, 2004; revised and accepted November 22, 2004.
Reprint requests: Dr. Ashok Agarwal, Professor of Surgery and Director, Center for Advanced Research in Human, Reproduction, Infertility, and Sexual Function, Glickman Urological Institute and Department of Obstetrics-Gynecology, The Cleveland Clinic Foundation, 9500 Euclid Avenue, Desk A19.1, Cleveland, OH 44195 (FAX: 216-445-6049; E-mail: agarwaa@ccf.org).

The process of cryopreservation increases the amount of apoptotic spermatozoa, which in turn decreases the success rate of ART. The binding of superparamagnetic annexin-V microbeads (ANMB) can effectively eliminate spermatozoa in early apoptotic stages from cryopreserved samples (12). Therefore, ANMB-negative spermatozoa may have higher survival potential after cryopreservation. The objective of our study was to determine if the inclusion of MACS in cryopreservation–thawing protocols improves sperm motility and the CSR.

MATERIALS AND METHODS

Sample Preparation

This study was approved by our institutional review board. Semen samples were collected from 10 healthy donors, and semen parameters exceeded the World Health Organization (13) reference ranges for the normal fertile population. To separate the predominantly mature spermatozoa, the liquefied semen was loaded onto a 55% and 80% discontinuous SupraSperm gradient (MediCult, Jyllinge, Denmark) and

centrifuged at $500 \times g$ for 20 minutes. The resulting 80% pellet (mature spermatozoa) was aspirated and resuspended in human tubal fluid media (HTF; Irvine Scientific, Santa Ana, CA).


The sperm cell suspension was divided into two separate fractions. The first was subjected to MACS followed by cryopreservation and thawing. The second was cryopreserved–thawed first and then subjected to MACS. Sperm motility was assessed in all fractions at each step of the experiment. The different steps of our experiment design are illustrated in Figure 1.

Isolation of Spermatozoa with Deteriorated Membranes by MACS

The sperm suspensions were passed through a magnetic field (MiniMACS; Miltenyi Biotec, Bergisch Gladbach, Germany), and the spermatozoa were classified as either ANMB-positive or ANMB-negative based on the binding of the microbeads to their outer surface (14).

FIGURE 1

Flow diagram of overall experiment design. Spermatozoa from same sample were subjected to cryopreservation–thawing before and after MACS. Number in parenthesis represents the aliquot number. MACS = magnetic activated cell separation; ANMB = annexin-V magnetic microbead; CSR = cryosurvival rate.



Said. Effect of MACS on sperm cryopreservation. Fertil Steril 2005.

Briefly, the washed spermatozoa were incubated with 100 μ L of ANMB at room temperature for 15 minutes, placed on top of the separation column containing iron balls, which was placed in a magnet. The apoptotic spermatozoa were retained in the separation column and labeled as ANMB-positive; the spermatozoa with intact membranes passed through the column and were labeled ANMB-negative. The power of the magnetic field was measured as 0.5 tesla between the poles of the magnet and up to 1.5 tesla within the iron globes of the column. After the column was removed from the magnetic field, the retained fraction was eluted using an annexin-binding buffer (15). The number of cells collected in each column exceeded 1×10^6 /mL.

Cryopreservation–Thawing Protocol

All specimens were cryopreserved using TEST-Yolk Buffer (TYB, 20% egg yolk and 12% glycerol; Irvine Scientific) (16). The TYB was added to the sperm samples at room temperature. An aliquot of the freezing medium equal to 25% of sperm sample volume was added to the specimen and gently mixed for 5 minutes using a Hema-Tek aliquot mixer (Miles Scientific, Elkhart, IN). This was repeated to give a final 1:1 (v/v) ratio of freezing medium to the sperm samples. Cryovials (1.5 mL; Corning, Pittsburg, PA) containing the specimens were placed in the freezer at -20°C for 8 minutes and thereafter in liquid nitrogen vapor at -80°C for 2 hours. The vials were finally transferred to liquid nitrogen tanks at -196°C . Twenty-four hours after the samples were frozen, a vial was removed and thawed by incubating it at 37°C for 20 minutes. Spermatozoa were washed and resuspended in HTF media immediately after thawing, and sperm motility was reassessed.

Statistical Analysis

Student's paired *t*-test was used to calculate the difference between samples. Hypothesis testing was two-tailed, and $P < .05$ was considered statistically significant. All values are given as mean \pm standard deviation (SD). All calculations were performed with Statistical 6.0 software (StatSoft, Tulsa, OK).

RESULTS

Raw semen samples collected from the donors had a sperm concentration of $89.5 \pm 22.3 \times 10^6$ /mL and percentage motility of 64.5 ± 6.43 . The percentage motility of the different aliquots are listed in Table 1. After MACS separation before freezing, the ANMB-negative sperm had statistically significantly higher motility values (76 ± 15.06 , $P = .03$) than the raw samples (64.5 ± 6.43) whereas the ANMB-positive sperm had statistically significantly lower values (41 ± 29.61 , $P < .0001$). The ANMB-negative sperm consistently had higher motility than the ANMB-positive sperm ($P = .006$).

The cryopreservation–thawing process statistically significantly decreased the motility of the spermatozoa that were not subjected to MACS separation compared with the pre-freeze control (20 ± 12.02 vs. 62.5 ± 20.45 , $P = .0001$). Similarly, the cryopreservation–thawing process decreased sperm motility in the ANMB-negative spermatozoa (38.5 ± 11.31 , $P = .01$) and in the ANMB-positive spermatozoa (4 ± 7 , $P < .0001$). However, the ANMB-negative sperm were the least affected by the cryopreservation–thawing process. The ANMB-negative sperm had higher motility following cryopreservation–thawing than the ANMB-positive sperm

TABLE 1

Motility values obtained from spermatozoa separated and not separated by magnetic-activated cell sorting (MACS).

Aliquot number	Aliquot content (n = 10)	MACS timing	CPT	Motility (%)
1	Raw semen	N/P	No	64.5 ± 6.43
2	Control (post density gradient)	N/P	No	62.5 ± 20.45
3	Control (post density gradient)	N/P	Yes	$20 \pm 12.02^{b,c}$
4	ANMB-negative	Prefreeze	No	76 ± 15.06^a
5	ANMB-positive	Prefreeze	No	41 ± 29.61^a
6	ANMB-negative	Prefreeze	Yes	38.5 ± 11.31^b
7	ANMB-positive	Prefreeze	Yes	$4 \pm 7^{b,c}$
8	ANMB-negative	Post-thaw	Yes	$2 \pm 2.58^{b,c}$
9	ANMB-positive	Post-thaw	Yes	$0.00 \pm 0.00^{b,c}$

Note: Values presented as mean \pm standard deviation; $P < .05$ considered statistically significant compared with ^araw semen, ^bnoncryopreserved control, and ^cANMB-negative spermatozoa separated by MACS and cryopreserved. N/P = MACS was not performed; ANMB = fraction separated by annexin-V magnetic beads; CPT = cryopreservation–thawing.

Said. Effect of MACS on sperm cryopreservation. Fertil Steril 2005.

($P < .0001$) and the sperm that were not separated by MACS ($P = .003$).

The use of TYB did not lead to any decrease in sperm motility. Motility values obtained after density gradient centrifugation were comparable with those obtained when TYB was added before cryopreservation in the samples that were not separated by MACS (62.5 ± 20.45 vs. 68.00 ± 10.33). When MACS was performed on the samples after the cryopreservation–thawing process, sperm motility was severely affected. Motility in the cryopreserved–thawed spermatozoa pre-MACS was statistically significantly higher than that in the post-MACS ANMB-positive cells (20.00 ± 12.02 vs. 0.00 ± 0.00 , $P = .0005$) and that of the ANMB-negative cells (20.00 ± 12.02 vs. 2.58 ± 0.81 , $P = .0005$).

The percentage of sperm that survived the cryopreservation process (percentage CSR) was calculated using the following formula: $100 \times \text{Post-thaw total motile sperm} / \text{Prefreeze total motile sperm}$. The CSR was highest in the ANMB-negative spermatozoa that were separated before freezing. This sperm aliquot had a statistically significantly higher CSR than the sperm cryopreserved without MACS (76.6 ± 59.75 vs. 30.29 ± 16.06 , $P = .04$) and the sperm that were ANMB-positive (76.6 ± 59.75 vs. 12.7 ± 31.19 , $P = .04$).

DISCUSSION

Despite various advances in cryopreservation methodology, the recovery rate of functional post-thaw spermatozoa remains unsatisfactory. Specifically, sperm motility significantly decreases after freezing (17). In our study, the cryopreservation–thawing process significantly decreased the percentage of motile spermatozoa. Sperm motility may have decreased because of a change in temperature and the formation and dissolution of ice in the extracellular environment (4). On the other hand, TYB may be excluded as a potentially damaging factor as sperm motility was maintained after it was added.

The cryopreservation–thawing process induces many changes in mammalian spermatozoa (18, 19). The lipid components of cell membranes tend to reorganize after being subjected to cooling, which may decrease the stability of the lipid bilayer (20, 21). In turn, sperm membrane integrity becomes impaired as PS translocates from the inner to the outer leaflet of the sperm plasma membrane (6, 22, 23), which is considered one of the early signs of apoptosis (7, 8). Annexin-V is a 35- to 36-kd phospholipid-binding protein that specifically binds to PS (6, 8, 9) and therefore can be used to detect early deleterious changes in the sperm plasma membrane that follow cryopreservation (24).

Superparamagnetic microbeads coupled with the use of specific antibodies can effectively separate cells. Based on the antibody used, leukocytes may be extracted from ejaculates, or an immature germ cell population may be separated from testicular tissue (25, 26). The beads may also be used

for immunomagnetic separation of membrane-intact and nonapoptotic spermatozoa (9, 12, 14, 27). The superparamagnetic annexin V-conjugated microbeads are able to eliminate spermatozoa with externalized PS (apoptotic cells) and disintegrated plasma membranes from cryopreserved semen samples (12, 27).

In the current study, we assessed the integration of MACS coupled with annexin V-conjugated microbeads in our cryopreservation protocol. The procedure delivers two sperm fractions: ANMB-positive (labeled apoptotic spermatozoa) and ANMB-negative (unlabeled with intact membranes). Before cryopreservation, ANMB-negative spermatozoa separated by MACS had the highest motility compared with the ANMB-positive and nonseparated spermatozoa. Following cryopreservation and thawing, ANMB-negative spermatozoa still had the highest motility and CSR compared with the other fractions. Therefore, it appears that the elimination of spermatozoa with early apoptotic changes positively affects motility and CSR after cryopreservation.

Sperm cryopreservation and thawing is associated with increased reactive oxygen species (ROS) production and decreased antioxidant levels (28, 29). In our earlier study, we noted that ROS levels have a positive correlation with the extent of apoptotic sperm (30). Therefore, the higher CSR in ANMB-negative sperm following cryopreservation–thawing could be related, at least in part, to the exclusion of the ROS-producing ANMB-positive sperm. In support, superoxide dismutase and catalase that act as selective scavengers for ROS were able to improve sperm recovery after cryopreservation–thawing (31).

In general, MACS is a feasible and safe method that may be used to produce a high-quality sperm fraction (9, 12, 14, 27). Although the separation columns and their magnetic field do not exert any detectable effect on the spermatozoa (12), sperm motility decreased dramatically in our current study when MACS was performed on cryopreserved–thawed samples. Because the deleterious effect of MACS was manifested only in the cryopreserved–thawed samples and not in the fresh samples, we therefore believe that this procedure can be integrated in cryopreservation protocols provided it is conducted on prefreeze specimens.

If used, MACS should be performed before cryopreservation. In our present study, sperm motility deteriorated when MACS was performed after cryopreservation, possibly because the cryopreservation procedures (e.g., centrifugation resuspension) made the cells vulnerable to damage. Moreover, ANMB-negative spermatozoa may still display a very early phase of PS translocation. In that case, only a limited number of beads would bind to the sperm—too few to be retained in the column.

Cryopreservation of human sperm is a fundamental tool for the preservation of male fertility. However, the process of cryopreservation impairs sperm fertility. Separating a distinctive population of nonapoptotic spermatozoa with in-

tact membranes before subjecting it to the cryopreservation–thawing process may optimize the outcomes. Our findings suggest that MACS coupled with ANMB not only can be used to perform this separation but can enhance sperm motility and CSR following cryopreservation.

REFERENCES

- Hovatta O. Cryobiology of ovarian and testicular tissue. *Best Pract Res Clin Obstet Gynaecol* 2003;17:331–42.
- Agarwal A, Ranganathan P, Kattal N, Pasqualotto F, Hallak J, Khayal S, et al. Fertility after cancer: a prospective review of assisted reproductive outcome with banked semen specimens. *Fertil Steril* 2004;81:342–8.
- Anger JT, Gilbert BR, Goldstein M. Cryopreservation of sperm: indications, methods and results. *J Urol* 2003;170:1079–84.
- Watson PF. The causes of reduced fertility with cryopreserved semen. *Anim Reprod Sci* 2000;60–61:481–92.
- Anzar M, He L, Buhr MM, Kroetsch TG, Pauls KP. Sperm apoptosis in fresh and cryopreserved bull semen detected by flow cytometry and its relationship with fertility. *Biol Reprod* 2002;66:354–60.
- Glander HJ, Schaller J. Binding of annexin V to plasma membranes of human spermatozoa: a rapid assay for detection of membrane changes after cryostorage. *Mol Hum Reprod* 1999;5:109–15.
- Vermes I, Haanen C, Steffens-Nakken H, Reutelingsperger C. A novel assay for apoptosis. Flow cytometric detection of phosphatidylserine expression on early apoptotic cells using fluorescein labelled Annexin V. *J Immunol Methods* 1995;184:39–51.
- Martin SJ, Reutelingsperger CP, McGahon AJ, Rader JA, van Schie RC, LaFace DM, et al. Early redistribution of plasma membrane phosphatidylserine is a general feature of apoptosis regardless of the initiating stimulus: inhibition by overexpression of Bcl-2 and Abl. *J Exp Med* 1995;182:1545–56.
- Glander HJ, Schiller J, Suss R, Paasch U, Grunewald S, Arnold J. Deterioration of spermatozoal plasma membrane is associated with an increase of sperm lyso-phosphatidylcholines. *Andrologia* 2002;34:360–6.
- Miltenyi S, Muller W, Weichel W, Radbruch A. High gradient magnetic cell separation with MACS. *Cytometry* 1990;11:231–8.
- von Schonfeldt V, Krishnamurthy H, Foppiani L, Schlatt S. Magnetic cell sorting is a fast and effective method of enriching viable spermatogonia from Djungarian hamster, mouse, and marmoset monkey testes. *Biol Reprod* 1999;61:582–9.
- Grunewald S, Paasch U, Glander HJ. Enrichment of non-apoptotic human spermatozoa after cryopreservation by immunomagnetic cell sorting. *Cell Tissue Bank* 2001;2:127–33.
- World Health Organization Laboratory manual for the examination of human semen and sperm-cervical mucus interaction. 4th ed. New York: Cambridge University Press, 1999.
- Paasch U, Grunewald S, Fitzl G, Glander HJ. Deterioration of plasma membrane is associated with activated caspases in human spermatozoa. *J Androl* 2003;24:246–52.
- Margolis LB, Namiot VA, Kljukin LM. Magnetoliposomes: another principle of cell sorting. *Biochim Biophys Acta* 1983;735:193–5.
- Hallak J, Sharma R, Wellstead C, Agarwal A. Cryopreservation of human spermatozoa: comparison of TEST-yolk buffer and glycerol. *Int J Fertil Womens Med* 2000;45:38–42.
- Donnelly ET, Steele EK, McClure N, Lewis SE. Assessment of DNA integrity and morphology of ejaculated spermatozoa from fertile and infertile men before and after cryopreservation. *Hum Reprod* 2001;16:1191–9.
- Medeiros CM, Forell F, Oliveira AT, Rodrigues JL. Current status of sperm cryopreservation: why isn't it better? *Theriogenology* 2002;57:327–44.
- Martin G, Sabido O, Durand P, Levy R. Cryopreservation induces an apoptosis-like mechanism in bull sperm. *Biol Reprod* 2004;71:28–37.
- Schiller J, Arnold J, Glander HJ, Arnold K. Lipid analysis of human spermatozoa and seminal plasma by MALDI-TOF mass spectrometry and NMR spectroscopy—effects of freezing and thawing. *Chem Phys Lipids* 2000;106:145–56.
- Fadok VA, Savill JS, Haslett C, Bratton DL, Doherty DE, Campbell PA, et al. Different populations of macrophages use either the vitronectin receptor or the phosphatidylserine receptor to recognize and remove apoptotic cells. *J Immunol* 1992;149:4029–35.
- Schuffner A, Morshedi M, Oehninger S. Cryopreservation of fractionated, highly motile human spermatozoa: effect on membrane phosphatidylserine externalization and lipid peroxidation. *Hum Reprod* 2001;16:2148–53.
- Duru NK, Morshedi MS, Schuffner A, Oehninger S. Cryopreservation–thawing of fractionated human spermatozoa is associated with membrane phosphatidylserine externalization and not DNA fragmentation. *J Androl* 2001;22:646–51.
- Pena FJ, Johannisson A, Wallgren M, Rodriguez-Martinez H. Assessment of fresh and frozen-thawed boar semen using an annexin-V assay: a new method of evaluating sperm membrane integrity. *Theriogenology* 2003;60:677–89.
- Hipler UC, Schreiber G, Wollina U. Reactive oxygen species in human semen: investigations and measurements. *Arch Androl* 1998;40:67–78.
- van der Wee KS, Johnson EW, Dirami G, Dym TM, Hofmann MC. Immunomagnetic isolation and long-term culture of mouse type A spermatogonia. *J Androl* 2001;22:696–704.
- Paasch U, Grunewald S, Agarwal A, Glandera HJ. Activation pattern of caspases in human spermatozoa. *Fertil Steril* 2004;81(Suppl 1):802–9.
- Chatterjee S, Gagnon C. Production of reactive oxygen species by spermatozoa undergoing cooling, freezing, and thawing. *Mol Reprod Dev* 2001;59:451–8.
- Bilodeau JF, Chatterjee S, Sirard MA, Gagnon C. Levels of antioxidant defenses are decreased in bovine spermatozoa after a cycle of freezing and thawing. *Mol Reprod Dev* 2000;55:282–8.
- Moustafa MH, Sharma RK, Thornton J, Mascha E, Abdel-Hafez MA, Thomas AJ Jr, et al. Relationship between ROS production, apoptosis and DNA denaturation in spermatozoa from patients examined for infertility. *Hum Reprod* 2004;19:129–38.
- Rossi T, Mazzilli F, Delfino M, Dondero F. Improved human sperm recovery using superoxide dismutase and catalase supplementation in semen cryopreservation procedure. *Cell Tissue Bank* 2001;2:9–13.